

090430F 0033.883.2100

PRESCRIPTION ANIMAL REMEDYKEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**LAURABOLIN® 25 mg/mL**

ANABOLIC STEROID FOR INTRAMUSCULAR OR SUBCUTANEOUS INJECTION

LAURABOLIN® 50 mg/mL

ANABOLIC STEROID FOR INTRAMUSCULAR OR SUBCUTANEOUS INJECTION

ACTIVE CONSTITUENT:
NANDROLONE LAURATE

For long term anabolic therapy in Horses, Dogs and Cats.

MODE OF ACTION:

LAURABOLIN® contains the highly active anabolic steroid 19 –norandrostenolone (Nandrolone) laurate in oil solution.

NANDROLIN® (Nandrolone Phenylpropionate B.P.) was the first anabolic agent to be made available with negligible androgenic activity for veterinary use and has gained wide acceptance in clinical practice. LAURABOLIN® (Nandrolone laurate) is an advanced development, aimed at providing for those conditions where fewer injections at longer intervals are desirable.

Thus where NANDROLIN® provides for short and medium term therapy, LAURABOLIN® provides for long term therapy. A single injection is sufficient to maintain continuous high level anabolic activity for 4-6 weeks. Where the acute stage of the condition has been controlled; the problem to be considered is that of convalescence. When the animal is established on the path to recovery, long term anabolic therapy using LAURABOLIN® will ensure adequate nitrogen for tissue synthesis. In addition, calcium and phosphorus retention is promoted, whilst normal levels of sodium, potassium, and water content of tissues are encouraged and maintained.

As a result of these effects, muscle power and function improve.

The healing of wounds and fractures is accelerated and an improvement in appetite has been reported.

INDICATIONS:

LAURABOLIN® is indicated in all cases requiring long term, less frequent injections following effective short term anabolic therapy.

Convalescence:

Following surgical operations and febrile diseases.

Newborn Animals:

Frequently foals are born in a weak condition which will respond well to anabolic steroid therapy.

Geriatrics:

Treatment of old animals is commonly complicated by renal dysfunction.

LAURABOLIN® may be used in these cases with the knowledge that the prolonged anabolic activity will have a desirable effect on nephritis with or without uraemia.

Tendon and Bone Damage:

In these conditions seen in the horse the repair processes are accelerated using LAURABOLIN®. Fracture healing is more rapid due to the positive effect on calcium and phosphorus metabolism.

Racing and working animals:

To achieve optimal fitness in racing and working animals a three month course of therapy is recommended.



Showing – one injection 3-4 weeks before.

If used in performance animals, the regulations of the relevant authorities regarding medication should be observed.

DIRECTIONS FOR USE:**RESTRAINT:****DO NOT USE in food producing species of animal.****PRECAUTIONS:**

Androgens can induce masculinisation of a female foetus and may have a suppressive effect on sexual development in young male animals.

DOSAGE AND ADMINISTRATION:

Use the contents of the vial within 24 hours of initial broaching. Discard any unused portion.

Give once each month by subcutaneous or intramuscular injection.

Dose rate is 1 mg/kg bodyweight.

Dogs up to 50 mg monthly.

Cats up to 25 mg monthly.

Horses up to 200 mg monthly.

Foals up to 100 mg monthly.

In chronic conditions or prolonged convalescence, injections should be made at three week intervals. In mares allow 6 weeks from injection date to mating date.

TOXICITY: 19 nor-androstenolone (as the decanoate) has been administered to dogs at four times normal doses over periods of six months without clinical or histological abnormality becoming evident. Overdosage may cause symptoms of androgenic activity.

MEAT WITHHOLDING PERIOD (HORSES): NOT TO BE USED in horses intended for human consumption.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764766.

DISPOSAL:

Dispose of empty container by wrapping with paper and putting in garbage.

STORAGE:

Store below 25°C (Air conditioning). Protect from light.

APVMA Approval No.: 37215 LAURABOLIN® 25 mg/mL

APVMA Approval No.: 37219 LAURABOLIN® 50 mg/mL

090430F 1103

INTERVET AUSTRALIA PTY LIMITED

91-105 Harpin Street, BENDIGO EAST VIC 3550, Phone: (03) 5440 9888

NEW ZEALAND INFORMATION:

It is a legal requirement that this product is not used in any animal producing or intended to produce food for human consumption.

INTERVET LTD., 49 George Street, NEWMARKET, AUCKLAND, Phone: (09) 309 0600

N.Z. Registered pursuant to the ACVM Act 1997 No. A1414 and A4638.

Prescription Animal Remedy (P.A.R.) Class 2.

For use only by, in the presence of, or under the control of a veterinarian.

Registered to INTERVET LTD., 49 George Street, Newmarket, Auckland.

See www.nzfsa.govt.nz/acvm/ for registration conditions.